

Retrospective studies into the effectiveness of the treatment of burns through distance healing.

By dr. Jan van Hemert, MD, Neurologist

Introduction.

Suppose, the simple fact is true that an atom is empty for 99.999% and cannot touch anything on an atomic level, because of the knocking off effect of electrons, the notion distance seems more likely to be normal than abnormal. Moreover, if the healing process seems to be evolving on an atom level, we may conclude that every healing process must be taking place from a distance. This process, however, is taking place in the subconscious, but, on the other hand, conscious manipulation is very possible. Even though this healing process rather seems to be a self-healing process than a process, which is being influenced from the outside.

Curing or healing must, in this case actually be seen as a tempo decrease of the self-healing process. The way the mechanism of using medication, a placebo or mental-manipulation works on a biological level doesn't seem clear yet. However, observed changes within the macroscopic world of healing cannot be denied and, moreover, we can only conclude that something is going on without us being able to grasp what is going on a molecular level or subatomic level. If methods of explanation are based on a set of scientific principles (like every scientific system does), excluding various explanatory modalities, a number of phenomenon's will never be explained from the simple conclusion that these observations are excluded beforehand from that scientific system. However, the denial of an observation, which does not fit in or is excluded from a scientific set of values is un-academic behavior and inhibits the development of science through exclusion.

I The Treatment of Burns From a Distance.

The mental interaction on life organism, have been studied extensively for decades (1-42). The Sahara experiments study the nature of influence from a distance. Clues resulting from these experiments prove that intentions from these interactions run through the autonomic nervous system. This notion is based on the positive findings of the galvanic skin test while being tested. The GSR concludes that the information, which is in the process of being transmitted from the emitted person to the receiving person is received on an unconscious level. The receiving person, however, conveys that he experiences a conscious sensation (this can be experiencing tingles, loss of equilibrium, a pressing sensation to the stomach or other sensations) that he has received the signal through conscious awareness. (Sahara experiments in stage of publication).

The treatment of burns in question is about a purposeful intention from one person to the other. The effectiveness of this treatment can be easily checked, because, in this case it is a matter of a shortening of the time span in the experience of pain in comparison to the regular time span between the experience of pain by people suffering from first or second-degree burns. One may conclude from the relevant literature that in the case of first degree burns the pain usually continues for 24 hours and in the case of a second degree burn the pain usually lingers for 24 hours. This way the results of the healing process are very easily measured in the percentage of time of being freed of pain. In comparison with the normal pain duration.

One may conclude from the available research that pain as such is caused by damage of the skin due to heat. The disappearance of pain is indicative of the speed in which pain subsides, this holds true together with the visual perception of the healing of the skin. The latter was clearly indicated by both patients observed by the therapist. The faster a wound heals will as such prove biologically the effectiveness of the intentional influence of the method of treatment and seems, moreover, to be the determining factor in this process which speeds up the healing process instead of only the disappearance of pain.

Because of a biological change, which occurs due to external causes, classic physicists talk about transformation of energy from one form to another. They are trying to state that the notion of labor means either the change of energy or the consumption of energy, no matter whether this particular source of energy is known. The statement that subtle energy would be responsible for this effect is totally unfounded. Moreover, from a scientific point of view it is of no significance. It is evident, however, that energy is an important factor. The nature of this healing energy remains unclear, however.

II Purpose of this Investigation.

The purpose of this study in question is to investigate the extent of the time-span, which needs to be determined by the easing of the pain concerning a regular time-span, which corresponds with the pain caused by first and second degree burns. This reduction of the amount of time was caused by this particular influencing of an intentional mental activity from a distance in case of such burns applied by trained persons.

III Material and Method.

A group of 64 people, who were lately treated, have been interviewed recently. This group consisted of 12 children, 13 men, and 39 women. From this group of people 34 persons suffered from first-degree burns. First-degree burns are considered to be burns.

In which only the epidermis is being damaged. Normally the pain lingers for about 24 hours and the healing of these wounds (recovery process) sets in within several days without remaining skin defects. From the group mentioned above, 30 of them suffered from second-degree burns. In a second-degree burn the epidermis and part of the true skin is being damaged, this goes together with the forming of blisters and scars.

In the opposite case, the deeper second-degree burns are more painful up to 4 weeks before total recovery, when relieve of pain sets in. The deeper kind of second degree burning is less painful, but goes together with a bad healing process of the skin.

Under investigation:

- 1 The period of time between the causing of the burns and enlisting someone's help.
- 2 The period of time between treatment and the total absence of pain. On a sliding scale from 100 till 0 we were only interested in zero.
- 3 Remarks from patients and observers about the healing process
- 4 The time space between the time of occurrence of the effect of second-degree burns and the time, which passed between the causing of the burn and the enlisting of help.

The method of treatment consisted of asking for the name, address, and the location of the burn on the body. The patient was told the treatment by means of concentration administered by the therapist, who focuses on the burnt location performed during 5 minutes. The patient was subsequently told that result could be expected in the same space off time between the burn and the call for help. The patient was kindly requested to send an e-mail containing the results including the specific times.

IV Results.

The results of the study are as follows:

- A. Children are totally free of pain for $1\frac{3}{4}$ hours, this is regardless of the degree of burning and the time between off burning and call for help. Usually, the lessening of the pain sets in immediately. It is not necessary for the patient to be aware of his treatment.

B. The 34 first-degree burns are free of pain within average time off 1 hour and 6 minutes.

C. The 30 second-degree burns are free of pain within average time off 1½ hours.

D. If the treatment of first-degree burns was started the average time of healing was 1 hour and 6 minutes.

V Conclusions.

The treatment of first-degree burns showed a complete disappearance of pain within 1 hour to 6 minutes. This stands in contrast to 24 hours of pain without being treated. This indicates a speeding up of the self-healing process of factor 22 in time.

In case of second -degree burns the time the patients was free of pain was 1.5 hours. This stands in contrast to the four weeks whenever the patient wasn't being treated. This indicates a speeding up of the self -healing process of factor 450 in time.

VI Discussion.

Because healing probably takes place on an atomic or even subatomic subconscious level, every type of influencing takes place from a distance, this is because on the level electrons never touch each other because of their negative charge. Here there is probably a healing-field involved, which is not tied to time or space. Moreover, the force of such a field should show an equal division, because of which the concept of distance loses its meaning independently off the distance between therapist and patient.

When an intervention takes place, the conscious concentration on the burnt body part is striking. If different areas, which are also burnt aren't mentioned by the patient during the telephone conversation, lessening of pain doesn't set on at these spots. In this case the patient should call the therapist again and once the therapist concentrates on the forgotten area, it will still become free of pain.

Up till now, the application of this method always has result according to the statements from the treated patients. The only difference would be the time the total halt of the pain sets in.

The essential question raised concerning pain control, is whether the notion pain can be dissociated from the healing of the burn. In other words, does the pain disappear due to rapid healing of the burn, or should we consider the state of halt

of pain dissociated from the healing process? Apart from visual observations of changes of the burn, the author isn't aware of a clear parameter, which will make the speed of the healing process measurable. So, we can only describe the various observations. However, there must certainly be a change in the aspect of the wound. Surely, this doesn't mean that we should actually be able to observe it. The aspect of the wound after the burn is naturally totally different from the aspect after 4 weeks of observing a second-degree burn without therapy.

However, the author and the therapist were an eyewitness to the changing of a burn within two hours, where one could observe in the time span of two hours when the burn closed itself with a thin skin tissue.

So we observed a leakage together with the start of small blister, a greenish aspect in the center of a yellowish color, changing slowly into a pink coloring with red dots and the disappearing of the hard red disk around the wound. In the mean time the wound was covered with a thin layer of skin accompanied with the easing of the pain. A dramatic observation by the therapist had taken place already, as she was able to actually witness the healing of a burn.

Usually, she heals from a distance, so she isn't able to observe any direct observation. She describes the same changes of the burn as is described above by the author. We are overall dependent on the descriptions of the patients.

So twenty patients out of 64 interviewed mention a direct visual healing of their wounds.

The obvious conclusion is that the alleviating of the pain is an attendant phenomenon of a faster healing process, which is going on and not an imagined psychological manifestation. So, it may be concluded, that the applied treatment may be considered to be speeding up the process of self-healing.

The gradual process of undulation of decreasing of pain while healing, which has been described by several patients could point into the direction of a biological factor, which may possibly biochemically determined. It might be interesting in case of future research to look for this possible biochemical link whatever it will be.

A Shortening of the time span of healing, could turn out to be of life saving importance in case of serious burns of more than 30% of the affected skin. The healing effect could surpass the speed of losing the serum caused by the burn.

It might also turn out to be interesting to apply the effect of intentional treatment on culture of skin tissue in the laboratory in order to find out whether the increase of growth can be influenced in a positive manner.

If there were biological parameters available of the self-healing effect, one could investigate which way and where the described method intervenes with the biological aspects of the cure.

If various therapists would apply this method to new patients, who suffer from various types of burns, they could be treated by these alternative ways, and, moreover, they could be treated at hospitals and specialized centers. This will prevent not only much suffering, but will also turn out to become an important money-saver. If a no cure no pay type of contract with this kind of therapy could be concluded, the risk for insurance companies may be driven back even further. It is our intention to supervise the entire course of treatment through research. This will enable a thorough evaluation of the entire process to take place.

VII Summary.

The application of the investigated method of healing from a distance shows an extraordinary high outcome of recovery from various types of burns. The shortening of the time of curing till 450 times of especially second-degree burns, includes the elaborate application to become widely applied to by centers treating patients suffering from burns.

References.

- 1&2) Barry, J.1967. Telepathy and plethysmography. *Revue Metapsychique* 6: 56-74
1968. General and comparative study of the psychokinetic effect on a fungus culture. *JP* 32:237-243
- 3) Beloff, J.1974. ESP: The search for a physiological index. *JSPR* 47: 401-420.
- 4&5) Benor, D.J. 1990. Survey of spiritual healing research. *Complementary Medical Research* 4: 9-33.
- 6) 1993. *Healing research*. Munich: Helix Verlag GmbH.
- 7) Beutler, J.J, J.T.M.Atteveld, et al. 1988 Paranormal healing and hypertension. *British Medical Journal* 296: 1491-1494.
- 8,9,10,11,12,13) Braud, W.G.1981.Psi performance and autonomic nervous system activity. *JASPR* 75: 1-35
1990.Distant mental influence of rate of hemolysis of human red blood cells. *JASPR* 84: 1- 24.
1993. on the use of living target systems in distant mental influence research. In *PSI research methodology: A re- examination*, edited by L.Coly. New York: Parapsychology Foundation and M.J. Schlitz 1983
Psychokinetic influence on electrodermal activity *JP* 47: 95-119.
1989. A methodology for the objective study of transpersonal imagery.*JSE* 3: 43-63.

1991. Consciousness interactions with remote biological systems:
Anomalous intentionality effects. SE2 (1): 1-46

- 14) Brier, R. 1969. PK on a bio-electrical system. JP 33: 187-205.
- 15) Campbell, A. 1968. Treatment of tumours by PK. JSPR 46: 428.
- 16) Delanoy, D., and S. Sha. 1994 Cognitive and psychological PSI responses to remote positive and neutral emotional states. In Proceedings of presented papers, 37th Annual Parapsychological Association Convention, edited by D.J. Bierman, 128-137. Fairhaven, MA: Parapsychological Association.
- 17) Dossey L., 1993. Healing words. San Francisco: Harper San Francisco.
- 18) Duane, T.D., and Behrendt. R. Extrasensory electroencephalographic induction between identical twins. Science 150: 367.
- 19) Gardner, R. 1983 Miracles of healing in Anglo-Celtic Northumbria as recorded by the Venerable Bede and his contemporaries: A reappraisal in the light of twentieth century experience. British Medical Journal 287: 1927-1933.
- 20&21) Grad, B. 1963. A telekinetic effect on plant growth. International JP 5: 117 - 134.
1965. Some biological effects of laying-on of hands: A review of experiments with animals and plants. JASPR 59: 95-127.
Cadoret R.J. and Paul G.K. 1961 The influence of an unorthodox method of treatment on wound healing in mice. International JP 3: 5-24
- 22) Grinberg-Zilberbaum, J., Delaflor M., Arellano M.E.S., Guevara M.A. and Perez M. 1992 Human communications and the electrophysiological activity of the brain. SE3 (3): 25-44.
- 23&24) Haraldsson, E. and Thorsteinsson T. 1973. Psychokinetic effects on yeast: An exploratory experiment. In RIP 1972, edited by W.G. Roll, R.L. Morris, and J.O. Morris, 20-21. Metuchen, NJ: Scarecrow Press.
- 25) Kuang, Ankun K., et al. 1986. Long-term observation on Qigong in prevention of stroke—Follow-up of 244 hypertensive patients for 18-22 years. Journal of Traditional Chinese Medicine 6 (4): 235-238.
- 26) May, E.C., Targ R. And Puthoff H.E. 1979. EEG correlates to remote light flashes under conditions of sensory shielding. In Mind at large, edited by C.T. Tart, H.E. Puthoff, and R. Targ, 127-136. New York: Praeger.

- 27) Miller, R.N. 1982 Study of remote mental healing. *Medical Hypotheses* 8:481 - 490
- 28) Morris R.L. 1977. Parapsychology, biology and anpsi. In *Handbook of parapsychology*, edited by B.B. Wolman, 687-716. New York: Van Nostrand Reinhold.
- 29&30) Nash, C.B. 1982 Psychokinetic control of bacterial growth. *JSPR* 51: 217-221. 1984. Test of psychokinetic control of bacterial mutation. *JASPR* 78: 145-152. and Nash C.S. 1967. The effect of paranormally conditioned solution on yeast fermentation. *JP* 31: 314.
- 31) Otani, S. 1955. Relations of mental set and change of skin resistance to ESP score. *JP* 19: 164-170.
- 32) Pleass, C.M, and Dey, D. 1990. Conditions that appear to favor extrasensory interactions between homo sapiens and microbes. *JSE4*: 213-231.
- 33) Radin, D.I., Taylor, R.D., and Braud, W. 1995 remote mental influence of human electrodermal activity: A pilot replication. *EJP11*: 19-34.
- 34) Randall, J.L. 1970. An attempt to detect psi effects with protozoa. *JSPR* 45
- 35) Rebman, J.M., Radin, D.I., Hapke R.A. and Gaughan, K. 1996. Remote influence of the autonomic nervous system by focused intention. In *Proceedings of Presented Papers, 39 th Annual Parapsychologic Association Convention*, edited by E.C. May, 133-148. Fairhaven, MA: Parapsychological Association.
- 36) Rebman, J.M., Wezelman, R., Radin D.I., Stevens P., Hapke, R. and Gaughan K. 1995. Remote influence of human physiology by a ritual healing technique. *SE* 6: 111-134.
- 37) Richmond, N. 1952. Two series of pk tests on paramecia. *JSPR* 36: 577-578.
- 38) Schouten, S.A. 1976 Autonomic Psychophysiological reactions to sensory and emotive stimuli in a psi experiment. *EJP* 1: 72-78.
- 39) Snel, F. Hol, P.R. 1983. Psychokinesis experiments in casein induced amyloidosis of the hamster. *EJP* 5 (1) 51-76.
- 40) Solfvin, G.F. 1982. PSI expectancy effects in psychic healing studies with malarial mice. *EJP* 4 (2): 160- 197.

- 41) Warren, C. McDonough, B.E., and Don, N.S. 1992. Event-related brain potential changes in a psi task. JP 56: 1-30.
- 42) Watkins, G.K. and Watkins A.M. 1971. Possible PK influence on the resuscitation of anesthetized mice. JP 35 (4): 257-272.

Author.

Dr. Jan van Hemert, Ph.D.

Jan van Hemert founder of the Teaching Tree University, was born in 1939 in the city of Rotterdam in the Netherlands. After finalizing his studies and medical specialization in Neurology and Neuro-Physiology, he became affiliated at the "Vrije University" as trainer of medical doctors and co-assistants. In addition, he was associated to the "Boerhave" method at the University of Leiden. Prof. Dr. Van Hemert was a successful medical director of several clinics based in Haarlem, the Netherlands. His interests however, went beyond neurology and he conducted research in the fields of Spasm, sickness of Parkinson, Epilepsy, Psychopharmacology, Stammer, Hyperventilation and condition improvement. He also became an expert in the areas of Chiropractic therapy, Neutral therapy, Auricle Therapy en Geo-biology. His school developed a new variety of the application of the "Lecher Antenna" and is right now in the process of developing a telegnostic device to establish quicker diagnosis in a non-invasive way with the cooperation of the University of Palermo in Italy. Jan established a research center and laboratory in his place of residence, being Egypt. His present research is mainly focused on the applications of holistic medicine and he is currently writing a manual for theoretical medicine based on the quantum theory and the intentionality within medicine. Jan van Hemert is also part of a group international medical doctors who are active in the campus medicus of the American Hospital in Rome.